

Know the basics

Being prepared and knowing the signs



Parents are usually good at noticing when something is wrong. However, it is normal to worry that you won't recognise the signs that your baby is unwell. Trust your instincts, you know your baby best.

Learn how to spot the signs of serious illness and how to cope if an accident happens. If you know the basics and you are prepared, you will find it easier to cope.

Keep a small supply of useful medicines in a **locked cabinet** or **somewhere up high** where a child cannot reach them. See box on the right, for things to have at home just in case. **Make sure you've got the right strength of medicine for the age of your child**, always follow instructions carefully and check use by dates. Read the label carefully. **Do not give aspirin to children under 16.**

Find out about CPR (resuscitation) before a possible emergency, visit www.redcrossfirstaidtraining.co.uk

If your baby seems to have a serious illness get medical help straight away.

Pharmacist says

Keep a small supply of useful medicines. Include things like:

- Thermometer
- Plasters
- Liquid painkillers (e.g. **sugar-free** paracetamol or ibuprofen)
- Barrier cream

Paracetamol and ibuprofen

Consider using either **sugar-free** paracetamol or ibuprofen for children with a fever who appear distressed (as a general rule a temperature of over **37.5 C (99.5 F)**, as these can help to reduce fever and distress. Treat them with either paracetamol **OR** ibuprofen in the first instance. It can take up to an hour for either of them to work. Paracetamol and ibuprofen should **NOT** be given together at the same time. However, if your child remains distressed before the next dose is due, then you may want to try a dose of the other medicine. Aspirin should not be given to children under 16 years of age.